



On Unitary Quasi-Equivalence and w-Hyponormal Operators In Hilbert Spaces

Kennedy Kibe Karanu ^{a*}, Zachary Kayiita Kaunda ^a
and Kinyanjui Jeremiah Ndung'u ^a

^aDepartment of Pure and Applied Sciences, Kirinyaga University, P. O. Box 143-10300, Kerugoya, Kenya.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jamcs/2024/v39i71909>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118609>

Received: 22/04/2024

Accepted: 27/06/2024

Published: 04/07/2024

Original Research Article

Abstract

In this paper, the properties of unitary quasi-equivalence on the class of w-hyponormal operators are presented using the Aluthge transform and polar decomposition property. We show that for any two unitary quasi-equivalent operators, F and G , if one is w-hyponormal, then the other operator is also w-hyponormal. This result also holds for p-hyponormal and log-hyponormal operators.

*Corresponding author: E-mail: kennedykibe34@gmail.com;

Cite as: Karanu, Kennedy Kibe, Zachary Kayiita Kaunda, and Kinyanjui Jeremiah Ndung'u. 2024. "On Unitary Quasi-Equivalence and W-Hyponormal Operators In Hilbert Spaces". Journal of Advances in Mathematics and Computer Science 39 (7):27-33. <https://doi.org/10.9734/jamcs/2024/v39i71909>.

Keywords: Hilbert space; unitary quasi-equivalence; w -hyponormal; p -hyponormal; log-hyponormal.

2010 Mathematics Subject Classification: 53C25; 83C05; 57N16.

1 Introduction

The Hilbert space, represented by H , is an inner product space that is complete [1]. The set of bounded linear operators in Hilbert spaces is denoted as $B(H)$. A mapping $F \in B(H)$ is an operator if it preserves the underlying structural properties of the space. An operator $F \in B(H)$ is hyponormal if $F^*F \geq FF^*$. An operator $F \in B(H)$ is said to have a polar decomposition if $F = U|F|$ where U is unitary. Two operators F and $G \in B(H)$ are said to be unitary quasi-equivalent if there exists a unitary operator U such that $F^*F = UG^*GU^*$ and $FF^* = UGG^*U^*$ where F^* denotes the adjoint of an operator F . This concept of unitary quasi-equivalence was introduced by Kutkut [2] under the idea of nearly equivalent operators by Othman [3]. Since then, various researchers have explored its properties on different operators in Hilbert space. Nzimbi and Wanyonyi [4] noted that unitary quasi-equivalence preserves the normality, hyponormality, and posinormality of operators. Lilian et al.[5] further established that unitary quasi-equivalence preserves the isometry, co-isometry, and partial isometry of an operator. Luketero and Khalagai [6] discovered that the operator is an equivalent relation. Luketero and Khalagai [6] also related unitary quasi-equivalence with other classes of equivalence relation, particularly unitary equivalence, and demonstrated that unitary quasi-equivalence implies unitary equivalence but the vice versa is not always true unless the operators are similar normal. Thereafter, Kikete et al. [7] introduced the concept of (n, m) -hyponormal and (n, m) -quasi-equivalence and proved it to be an equivalence relation. Kikete et al. [7] also proved that (n, m) -hyponormal preserves the (n, m) -quasi-equivalence of an operator. That is, if two operators are (n, m) -unitary quasi-equivalence and one of them is (n, m) -hyponormal then the other must be (n, m) -hyponormal. However, similar results on the class of w -hyponormal, p -hyponormal, and log-hyponormal operators have not been shown. Thus, this study focused on determining the properties of unitary quasi-equivalence on log-hyponormal, p -hyponormal, and w -hyponormal operators.

2 Definitions

Definition 2.1: hyponormal Operator [8]. An operator $F \in B(H)$ is hyponormal if $F^*F \geq FF^*$.

Definition 2.2: w -hyponormal operator [9]. An operator F is said to be w -hyponormal if it satisfies the condition $|\tilde{F}| \geq |F| \geq |\tilde{F}^*|$.

Definition 2.3: log-hyponormal [10], [11]. An invertible operator F is called log-hyponormal if $\log(F^*F) \geq \log(FF^*)$.

Definition 2.4: p -hyponormal [12], [13]. An operator $F \in B(H)$ is said to be p -hyponormal if $(F^*F)^p - (FF^*)^p \geq 0$ for $0 < p \leq 1$.

Definition 2.5: Aluthge Transform [13]. The first Aluthge transform of F is defined as $\tilde{F} = |F|^{1/2}U|F|^{1/2}$ where $F = U|F|$, while the second transform of F is defined as $\tilde{\tilde{F}} = |\tilde{F}|^{1/2}\tilde{U}|\tilde{F}|^{1/2}$.

Definition 2.6: Unitary Quasi-equivalence Operator [5]. F and $G \in B(H)$ are said to be unitary quasi-equivalent if there exists a unitary operator U such that:

$$F^*F = UG^*GU^* \quad \text{and} \quad FF^* = UGG^*U^*$$

From the definition above, the clas of w -hyponormal p -hyponormal and log-hyponormal operators relates as follows;

$$\begin{aligned} & \text{Hyponormal} \subset \text{Log-hyponormal} \subset w\text{-Hyponormal} \subset \text{Paranormal} \subset K\text{-paranormal} \\ & \text{Hyponormal} \subset p\text{-Hyponormal} (0 < p < 1) \subset w\text{-Hyponormal} \subset \text{Paranormal} \subset K\text{-paranormal}[14] \end{aligned}$$

3 Main Results

Lemma 3.1 [4] Let $F, G \in B(H)$ be unitarily quasi-equivalent. Then F is hyponormal if and only if G is hyponormal.

Lemma 3.2 [6] Let F be a hyponormal operator and G be another operator such that:

1. $F = UGU^*$ where U is an isometry, or
2. $F = U^*GU$ where U is a co-isometry.

Then G is also hyponormal.

Remark: From preceding lemmas it is evident that hyponormal operators are invariant under unitary quasi-equivalence, isometric and co-isometric transformations. To extend these result, the study aimed to establish the following results.

Theorem 3.3: If $F, G \in B(H)$ are unitary quasi-equivalent operators and F is a log-Hyponormal operator, then G is also log-Hyponormal.

Proof. Since F is unitary quasi-equivalent to G , then

$$F^*F = UG^*GU^* \tag{1}$$

$$FF^* = UGG^*U^* \tag{2}$$

But F is log-Hyponormal thus by definition

$$\log(F^*F) \geq \log(FF^*) \tag{3}$$

Substituting equations (1) and (2) into (3)

$$\log(UG^*GU^*) \geq \log(UGG^*U^*) \tag{4}$$

Pre-multiplying and post-multiplying UG^*GU^* and UGG^*U^* in equation (4) with U^* and U on the left and on the right, results to;

$$\log(U^*UG^*GU^*U) \geq \log(U^*UGG^*U^*U) \tag{5}$$

But

$$U^*U = UU^* = I$$

Thus (5) becomes

$$\begin{aligned} \log(IG^*GI) &\geq \log(IGG^*I) \\ \log(G^*G) &\geq \log(GG^*) \end{aligned} \tag{6}$$

Thus by definition G is a log-hyponormal operator. □

Theorem 3.4: Suppose $F, G \in B(H)$ are unitary quasi-equivalent operators. If F is a p -hyponormal operator, then G is also a p -hyponormal operator.

Proof. Since $F \overset{u.q.e}{\approx} G$ then we have that;

$$F^*F = UG^*GU^* \tag{7}$$

$$FF^* = UGG^*U^* \tag{8}$$

However, F is p -hyponormal thus by definition:

$$(F^*F)^p \geq (FF^*)^p \tag{9}$$

Replacing equations (7) and (8) into equation (9):

$$(UG^*GU^*)^p \geq (UGG^*U^*)^p \tag{10}$$

By the polar decomposition property:

$$G = U|G| \text{ where } U \text{ is a unitary operator.} \tag{11}$$

$$G^* = |G^*|U^* \tag{12}$$

Substituting equations (11) and (12) into equation (10):

$$(U|G^*|U^*U|G|U^*)^p \geq (UU|G||G^*|U^*U^*)^p \tag{13}$$

From definition, equation (13) becomes:

$$U((|G^*|U^*U|G|)^p)U^* \geq U((U|G||G^*|U^*)^p)U^* \tag{14}$$

Post-multiplying and pre-multiplying both sides of equation (14) with U^* on the left and U on the right;

$$U^*U((|G^*|U^*U|G|)^p)U^*U \geq U^*U((U|G||G^*|U^*)^p)U^*U \tag{15}$$

But $UU^* = U^*U = I$. Replacing the identity in equation (15):

$$\begin{aligned} I((|G^*|U^*U|G|)^p)I &\geq I((U|G||G^*|U^*)^p)I \\ ((|G^*|U^*U|G|)^p) &\geq ((U|G||G^*|U^*)^p) \end{aligned} \tag{16}$$

But from equations (11) and (12), (16) simplifies to:

$$(G^*G)^p \geq (GG^*)^p \tag{17}$$

And hence from (17) G is a p -hyponormal operator. \square

Theorem 3.5: If F and G are projection and unitary equivalent operators, and F is a w -hyponormal operator then G is also w hyponormal.

Proof. From hypothesis $F \stackrel{u.q.e}{\approx} G$ implying that:

$$F^*F = UG^*GU^*$$

And

$$FF^* = UGG^*U^*$$

Also F is w -hyponormal which means that

$$|\tilde{F}| \geq |F| \geq |\tilde{F}^*| \tag{18}$$

Where:

$$F = U|F| \tag{19}$$

$$|\tilde{F}| = |F|^{1/2}U|F|^{1/2} \tag{20}$$

$$|\tilde{F}^*| = |F^*|^{1/2}U^*|F^*|^{1/2} \tag{21}$$

Substituting equations (19), (20), and (21) into equation (18) we get:

$$\||F|^{1/2}U|F|^{1/2}\| \geq \|U|F|\| \geq \| |F^*|^{1/2}U^*|F^*|^{1/2} \| \tag{22}$$

Since F is a projection,

$$F^* = UG^*U^*$$

And

$$F = UGU^*$$

Substituting F and F^* into equation (22) we get:

$$\| |UGU^*|^{1/2}U|UGU^*|^{1/2} \| \geq \|U|UGU^*|\| \geq \| |UG^*U^*|^{1/2}U^*|UG^*U^*|^{1/2} \| \tag{23}$$

By the polar decomposition property equation (21) can be rewritten as:

$$U(|G|^{1/2})U^*UU(|G|^{1/2})U^* \geq UU|G|U^* \geq U(|G^*|^{1/2})U^*U^*U(|G^*|^{1/2})U^* \tag{24}$$

Since U is unitary, then

$$U^*U = I \quad \text{where } I \text{ is the identity.}$$

Thus:

$$U(|G|^{1/2})U(|G|^{1/2})|U^* \geq UU||G||U^* \geq U(|G^*|^{1/2})U^*I(|G^*|^{1/2})|U^* \tag{25}$$

Pre-multiplying both sides of equation (25) by U^* on the left and post-multiplying U on the right side we get:

$$U^*U(|G|^{1/2}U|G|^{1/2})U^*U \geq U^*UU||G||U^*U \geq U^*U(|G^*|^{1/2}U^*I|G^*|^{1/2})U^*U$$

But

$$U^*U = I$$

$$I(|G|^{1/2}U|G|^{1/2})I \geq IU||G||I \geq I(|G^*|^{1/2}U^*I|G^*|^{1/2})I$$

Thus

$$\|(|G|^{1/2}U|G|^{1/2})\| \geq \|U||G|\| \geq \| (|G^*|^{1/2}U^*|G^*|^{1/2})\| \tag{26}$$

But

$$\begin{aligned}(|G|^{1/2}U|G|^{1/2}) &= |\tilde{G}| \\ U|G| &= T\end{aligned}$$

And

$$(|G^*|^{1/2}U^*|G^*|^{1/2}) = |\tilde{G}^*|$$

Thus equation (26) simplifies to

$$|\tilde{G}| \geq |G| \geq |\tilde{G}^*|$$

And hence by definition 2.2 G is also w -hyponormal. □

4 Conclusion

The preceding results has shown that unitary quasi-equivalence preserves the properties of w -hyponormal, log-hyponormal and p -hyponormal. That is for any two operators which are unitary quasi-equivalence and one of them being w -hyponormal, log-hyponormal or p -hyponormal then the other operators must also be w -hyponormal, log-hyponormal or p -hyponormal. These findings will be valuable for those using these operators in quantum mechanics to develop theoretical formulations. Also, the results will be significant in applied algebra and the study of differential operators for the calculation and differentiation of wave functions.

Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Halmos PR. A Hilbert space problem book. Springer Science & Business Media; 2012 Dec 6.
- [2] Mahmoud Kutkut. On quasiequivalent operators, Bull. Cal. Math. Soc. 1998;90:45-52.
- [3] Sadoon Ibrahim Othman. Nearly equivalent operators, Math. Bohem. 1996;212(2):133-141.
- [4] Nzimbi BM, Wanyonyi SL. On unitary quasi-equivalence of operators. International Journal of Mathematics and Its Applications. 2020;8(1):207-215.
- [5] Lilian A, Wabomba MS, Ndung'u KJ. On Unitary Quasi-equivalence and partial isometry operators in hilbert spaces. Journal of Advances in Mathematics and Computer Science. 2023 Oct 18;38(10):113-20.
- [6] Luketero SW, Khalagai JM. On unitary equivalence of some classes of operators in Hilbert spaces; 2020.
- [7] Kikete DW, Luketero SW, Mile JK, Wafula AW. A study on properties of $(n; m)$ -hyponormal operators. Asian Journal of Pure and Applied Mathematics. 2023 Jul 3;209-17.

- [8] Stampfli, Joseph. Hyponormal operators. Pacific J. Math. 1962;12(4):1453-1458.
- [9] Rashid MH. Some invariant subspaces for w -hyponormal operators. Linear and Multilinear Algebra. 2019 Jul 3;67(7):1460-70.
- [10] Tanahashi K. On log-hyponormal operators, Integr. Equat. Oper. Th. 1999;34:364372
- [11] Prasad T, Bachir A. Fuglede-putnam theorem for pw -hyponormal or class Y operators. Int. J. Oper. Problems Comput. Math. 2018 Mar 1;11(1).
- [12] Shagari MS. On normality conditions of p -hyponormal operators. J. Anal. Num. Theor. 2019;1-6.
- [13] Aluthge A. On p -hyponormal operators for $0 < p < 1$. Integral Equations and Operator Theory.1990 May;13:307-15.
- [14] Mutuku S. On Aluthge Transforms and Spectral Properties of different classes of operators in Hilbert spaces (Doctoral dissertation, University of Nairobi); 2020.

Disclaimer/Publisher's Note:The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/118609>