



Automorphism of Zero Divisor Graphs of Nilradicals of Commutative Finite Local Rings

Presley Kiplagat ^{a*}, Lao Hussein Mude ^a and Zachary Kaunda Kayiita ^a

^a Department of Pure and Applied Sciences, Kirinyaga University, P. O. Box 143-10300, Kerugoya, Kenya, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jamcs/2026/v41i22097>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://pr.sdiarticle5.com/review-history/152768>

Received: 29/11/2025

Published: 07/02/2026

Original Research Article

Abstract

A zero-divisor graph of a commutative ring R denoted as $\Gamma(R)$, is a graph whose vertices are the zero divisors of the ring. Any two distinct vertices of the graph are incident if and only if their product is zero. The zero-divisor graph associated with a commutative ring encodes deep algebraic information in a combinatorial framework. In this paper, we investigate the automorphism groups of zero-divisor graphs arising from the nonzero nilradical of finite local rings of the form \mathbb{Z}_{p^k} . By exploiting the natural p -adic valuation on nilpotent elements, we obtain a canonical stratification of the vertex set into valuation levels. This structure allows for a precise description of graph automorphisms as products of symmetric groups acting on valuation classes. The results provide a complete characterization of graph symmetries in this local setting and establish a foundational case for the broader theory of automorphisms of zero-divisor graphs over finite rings.

Keywords: Zero-divisor graph; nilradical; automorphism group; local ring; p-adic valuation.

*Corresponding author: E-mail: presleykiplagat@gmail.com;

Cite as: Kiplagat, Presley, Lao Hussein Mude, and Zachary Kaunda Kayiita. 2026. "Automorphism of Zero Divisor Graphs of Nilradicals of Commutative Finite Local Rings". *Journal of Advances in Mathematics and Computer Science* 41 (2):28-33. <https://doi.org/10.9734/jamcs/2026/v41i22097>.

2010 Mathematics Subject Classification: 53C25, 83C05, 57N16.

1 Introduction

Zero-divisor graphs have become an important interface between commutative algebra and graph theory since the seminal work of Anderson and Livingston (Anderson and Livingston, 1999). By encoding algebraic interactions among zero divisors as adjacency relations, these graphs enable the study of algebraic properties of rings via graph-theoretic invariants such as connectivity, diameter, girth, clique number, chromatic number, and binding number (Ali et al., 2024, 2025). A substantial body of literature has since developed around these themes, particularly for finite commutative rings and Artinian local rings (Beck, 1988; DeMeyer and Schneider, 2003). Much of the existing research has focused on the structural properties of $\Gamma(R)$ and its variants, including Beck's total graph (Beck, 1988). Results by Anderson and Livingston (1999); Akbari et al. (2003); Axtell and Stickles (2009), and others establish sharp bounds on diameter and girth, characterize completeness and multipartite structure, and relate these graph invariants to ideal-theoretic properties such as nilpotency, chain conditions on prime ideals, and the behavior of the Jacobson radical. In particular, for local rings, the structure of the nilradical plays a decisive role in determining the shape of the associated zero-divisor graph (Lucas, 2006; Sbarra and Zanardo, 2024). This role of the nilradical has also been emphasized in the work of Anderson and Badawi (2008), where annihilator-based partitions are shown to strongly influence graph structure. In contrast, comparatively little attention has been paid to the automorphism groups of zero-divisor graphs, especially those arising from nilradicals. While it is well understood that $\Gamma(R)$ often reflects ring-theoretic symmetries, explicit descriptions of $\text{Aut}(\Gamma(R))$ are known only for restricted classes of rings (DeMeyer and Schneider, 2003). Existing studies on graph automorphisms have largely focused on reduced rings, Gorenstein rings, or special constructions such as idealizations and direct products (Spiroff and Wickham, 2011). Even in the case of finite local rings, where $\Gamma(R)$ is finite and structurally constrained, the automorphism groups have not been systematically described in terms of intrinsic algebraic data, apart from specific families such as Galois rings (Mude et al., 2023; Onyango et al., 2016). This gap is particularly evident for principal ideal local rings such as \mathbb{Z}_{p^k} . Although the zero-divisor graphs of these rings are well understood with respect to diameter, girth, and completeness (Anderson and Livingston, 1999; Lucas, 2006), the precise manner in which p -adic valuation and nilpotency stratify the vertex set—and thereby restrict graph automorphisms—has not been fully articulated. More broadly, the extent to which valuation-theoretic data determines $\text{Aut}(\Gamma)$ for nilradicals remains unclear, for local rings cases. The purpose of this paper is to address this gap by providing an analysis of the automorphism groups of zero-divisor graphs arising from the nonzero nilradical of finite local rings of the form \mathbb{Z}_{p^k} . We show that the p -adic valuation induces a canonical stratification of the vertex set into invariant layers, each consisting of vertices with identical graph-theoretic neighborhoods. This stratification forces the automorphism group to decompose as a direct product of symmetric groups acting independently on valuation classes. Beyond yielding explicit formulas for $\text{Aut}(\Gamma)$, this approach clarifies the structural reason behind the observed symmetry: in local rings, valuation levels act as rigid combinatorial invariants. The results presented here form a foundational case that informs and motivates subsequent investigations of semilocal and mixed-exponent rings, where similar valuation-based mechanisms govern graph automorphisms but interact across multiple maximal ideals.

2 Algebraic Setting and Nilpotent Structure

Throughout this section, we consider the principal ideal local ring \mathbb{Z}_{p^k} , where p is a prime and $k \geq 2$. As a finite commutative local ring with a unique maximal ideal (p) , the structure of \mathbb{Z}_{p^k} is governed entirely by p -adic valuation. Understanding the precise form of nilpotent elements is essential for the study of zero-divisor graphs of nilradicals. Since annihilation conditions control adjacency in these graphs modulo p^k , the p -adic valuation of each vertex serves as a fundamental invariant.

Theorem 2.1. *Let $R = \mathbb{Z}_{p^k}$, where p is a prime and $k \geq 2$. Then an element $x \in R$ is nilpotent if and only if $p \mid x$. Moreover, every nonzero nilpotent element admits a unique representation $x = p^r a$, $1 \leq r \leq k-1$, $\gcd(a, p) = 1$.*

Proof. Since \mathbb{Z}_{p^k} is a finite commutative ring, its nilradical coincides with the intersection of all prime ideals. Because \mathbb{Z}_{p^k} is local with unique maximal ideal (p) , this intersection is precisely (p) . Equivalently, the nilradical of \mathbb{Z}_{p^k} is generated by

p , and an element is nilpotent if and only if it belongs to the ideal (p) . Let $x \in \mathbb{Z}_{p^k}$. If $p \nmid x$, then x is a unit in R , and hence no power of x can be zero. Therefore, x is not nilpotent. Conversely, if $p \mid x$, then $x \in (p)$, and since $(p)^k = (0)$ in \mathbb{Z}_{p^k} , it follows that some power of x must be zero. This proves that x is nilpotent if and only if $p \mid x$. Now suppose x is a nonzero nilpotent element. By the fundamental theorem of arithmetic, there exists a unique integer $r \geq 1$ such that $x = p^r a$, where a is an integer satisfying $\gcd(a, p) = 1$. Since $x \not\equiv 0 \pmod{p^k}$, we must have $r \leq k - 1$. This yields the stated representation. To verify nilpotency explicitly, observe that $x^m = p^{rm} a^m \equiv 0 \pmod{p^k}$ if and only if $rm \geq k$. Because $r \geq 1$, such an integer m always exists, confirming that x is nilpotent. Finally, the representation $x = p^r a$ is unique, since the p -adic valuation $v_p(x) = r$ is uniquely determined and the residue class of a modulo p^k is fixed once r is specified. \square

3 Canonical Decomposition of the Vertex Set

A central step in understanding the automorphism group of a zero-divisor graph is the identification of intrinsic partitions of the vertex set that are dictated by the underlying algebraic structure of the ring (Patil et al., 2022). For the ring \mathbb{Z}_{p^k} , the natural invariant governing this structure is the p -adic valuation. As shown in the preceding section, every nonzero nilpotent element admits a unique factorization of the form $p^r a$ with $1 \leq r \leq k - 1$ and $\gcd(a, p) = 1$. This representation immediately suggests a stratification of the vertex set according to valuation level. From a graph-theoretic perspective, such a stratification is not merely algebraically convenient: vertices with the same p -adic valuation exhibit identical annihilation behavior and hence share the same adjacency relations in the zero-divisor graph. Conversely, vertices with different valuations interact differently from other elements of the nilradical.

Theorem 3.1. *The vertex set of the zero-divisor graph $\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))$ admits the disjoint decomposition $V = \bigsqcup_{r=1}^{k-1} S_{p^r}$, $S_{p^r} = \{p^r a : \gcd(a, p) = 1\}$, with $|S_{p^r}| = p^{k-r-1}(p - 1)$.*

Proof. Let x be a vertex of $\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))$. By definition, x is a nonzero nilpotent element of \mathbb{Z}_{p^k} . From the characterization of nilpotent elements established earlier, there exists a unique integer r with $1 \leq r \leq k - 1$ and a unique residue class a satisfying $\gcd(a, p) = 1$ such that $x = p^r a$. This shows that every vertex belongs to at least one subset S_{p^r} . To see that the union is disjoint, suppose that $x \in S_{p^r} \cap S_{p^s}$ for some $r \neq s$. Then $x = p^r a = p^s b$ for some a, b coprime to p . Without loss of generality, assume $r < s$. Dividing both sides by p^r yields $a = p^{s-r} b$, which implies that $p \mid a$, contradicting the assumption $\gcd(a, p) = 1$. Hence each vertex belongs to exactly one valuation class S_{p^r} , and the decomposition is disjoint and exhaustive. We now compute the cardinality of each subset S_{p^r} . Fix an integer r with $1 \leq r \leq k - 1$. Distinct elements of S_{p^r} correspond precisely to distinct residue classes of a modulo p^{k-r} such that $\gcd(a, p) = 1$. Indeed, if $p^r a \equiv p^r a' \pmod{p^k}$, then $a \equiv a' \pmod{p^{k-r}}$. Thus, counting elements of S_{p^r} is equivalent to counting the units of the ring $\mathbb{Z}_{p^{k-r}}$. The number of units modulo p^{k-r} is given by Euler's totient function: $\phi(p^{k-r}) = p^{k-r} - p^{k-r-1} = p^{k-r-1}(p - 1)$. Therefore, $|S_{p^r}| = p^{k-r-1}(p - 1)$. \square

4 Adjacency Relations and Degree Invariants

Since adjacency is defined through multiplicative annihilation modulo p^k , it is natural to expect that the sum of valuations of two vertices plays a decisive role. Indeed, the product of two nilpotent elements vanishes modulo p^k precisely when the combined power of p in their factorization reaches or exceeds the modulus exponent. From a graph-theoretic standpoint, this observation has two important consequences. First, it yields a simple and explicit criterion for adjacency that depends only on valuation levels and not on unit coefficients. Second, it implies that all vertices within a fixed valuation class have identical neighborhoods and hence identical degrees. This uniformity is a crucial structural feature: degree is a graph invariant preserved by automorphisms, and thus valuation classes become rigid building blocks in the symmetry analysis of $\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))$.

Theorem 4.1. *Let $x = p^r a$ and $y = p^s b$ be distinct vertices of $\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))$. Then x is adjacent to y if and only if $r + s \geq k$. Consequently, all vertices in the same set S_{p^r} have identical degrees.*

Proof. By definition of the zero-divisor graph, two distinct vertices x and y are adjacent if and only if $xy \equiv 0 \pmod{p^k}$. Using the unique p -adic representations $x = p^r a$ and $y = p^s b$, where $\gcd(a, p) = \gcd(b, p) = 1$, we compute $xy = p^{r+s} ab$.

Since a and b are units modulo p , their product ab is also a unit modulo p . In particular, ab is not divisible by p , and therefore contributes no additional p -power to the factorization of xy . It follows that the highest power of p dividing xy is exactly p^{r+s} . Consequently, $xy \equiv 0 \pmod{p^k} \iff p^k \mid p^{r+s} \iff r+s \geq k$. This establishes the claimed adjacency criterion. We now examine the degree of a vertex $x = p^r a \in S_{p^r}$. By the criterion just proved, x is adjacent precisely to those vertices $y = p^s b$ for which $s \geq k - r$. Thus the neighborhood of x is given by $N(x) = \bigcup_{s=k-r}^{k-1} S_{p^s}$. Importantly, this description depends only on the valuation level r and not on the specific choice of the unit a . Since the cardinalities $|S_{p^s}|$ are fixed for each s , the degree of x is $\deg(x) = \sum_{s=k-r}^{k-1} |S_{p^s}|$, which is independent of the particular vertex chosen within S_{p^r} . Hence all vertices belonging to the same valuation class S_{p^r} have identical degrees. \square

5 Invariance of Valuation Levels

Since automorphisms preserve graph-theoretic invariants derived from adjacency, they cannot arbitrarily rearrange vertices across valuation classes without violating these invariants (Madhumitha and Naduvath, 2023). This observation suggests that valuation subsets should be fixed setwise by every graph automorphism. The result below formalizes this intuition and establishes valuation levels as rigid combinatorial strata that govern the symmetry structure of $\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))$.

Theorem 5.1. *Each valuation subset S_{p^r} is invariant under $\text{Aut}(\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k})))$.*

Proof. From Theorem 4.1, the degree of a vertex $x = p^r a$ depends only on its valuation level r . More precisely, all vertices in S_{p^r} share the same degree, while vertices belonging to distinct valuation subsets S_{p^r} and S_{p^s} with $r \neq s$ have different degrees. Let $\phi \in \text{Aut}(\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k})))$. Since ϕ is a graph automorphism, it preserves adjacency and therefore preserves vertex degrees. Hence, for any vertex $x \in S_{p^r}$, the image $\phi(x)$ must have the same degree as x . If $\phi(x)$ were to lie in a different valuation subset S_{p^s} with $s \neq r$, then $\deg(\phi(x)) = \deg(x)$ would contradict the fact that vertices in S_{p^r} and S_{p^s} have distinct degrees. Therefore, no such mapping is possible. It follows that $\phi(S_{p^r}) \subseteq S_{p^r}$ for all $r = 1, \dots, k-1$. Since ϕ is bijective, the inclusion must in fact be equality. Thus each valuation subset S_{p^r} is preserved setwise by every automorphism of the graph. This establishes the invariance of valuation levels under $\text{Aut}(\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k})))$. \square

6 Automorphism Group Decomposition

The preceding sections establish two fundamental structural facts about $\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))$: first, the vertex set decomposes into valuation-defined subsets S_{p^r} , and second, these subsets are invariant under graph automorphisms. Since all vertices in a fixed valuation class share identical adjacency relations, they are indistinguishable from a purely graph-theoretic perspective. This symmetry suggests that automorphisms act freely within valuation layers but are prohibited from mixing vertices across different layers.

Theorem 6.1. *Let $R = \mathbb{Z}_{p^k}$ with $k \geq 3$. Then $\text{Aut}(\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))) \cong \prod_{r=1}^{k-1} S_{p^{k-r-1}(p-1)}$.*

Proof. By Theorem 5.1, each valuation subset S_{p^r} is invariant under every graph automorphism. Hence, any automorphism ϕ restricts to a permutation of each S_{p^r} , yielding a homomorphism

$$\text{Aut}(\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))) \longrightarrow \prod_{r=1}^{k-1} \text{Sym}(S_{p^r}).$$

By Theorem 4.1, all vertices within a fixed subset S_{p^r} have identical neighborhoods. Consequently, any permutation of S_{p^r} preserves adjacency with all other vertices of the graph. It follows that every permutation of S_{p^r} extends to a graph automorphism that fixes all other valuation levels pointwise. This shows that the above homomorphism is surjective. Injectivity follows from the fact that an automorphism acting trivially on each S_{p^r} must fix every vertex of the graph and hence is the identity automorphism. Therefore, $\text{Aut}(\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))) \cong \prod_{r=1}^{k-1} \text{Sym}(S_{p^r})$. Finally, since $|S_{p^r}| = p^{k-r-1}(p-1)$ by Theorem 3.1, the result follows. \square

Remark 6.1. When $R = \mathbb{Z}_{p^2}$, every nonzero nilpotent element is of the form pa with $\gcd(a, p) = 1$, so the vertex set reduces to a single valuation class $V(\Gamma(\text{Nil}^*(\mathbb{Z}_{p^2}))) = S_p$, $|S_p| = p - 1$. For any two distinct vertices $x = pa$ and $y = pb$, we have $xy = p^2ab \equiv 0 \pmod{p^2}$, so every pair of distinct vertices is adjacent. Hence, $\Gamma(\text{Nil}^*(\mathbb{Z}_{p^2})) \cong K_{p-1}$, the complete graph on $p - 1$ vertices. In this situation, the valuation-based layering collapses entirely, and there are no degree or neighborhood distinctions among vertices. Consequently, every permutation of the vertex set preserves adjacency, and the automorphism group is the full symmetric group $\text{Aut}(\Gamma(\text{Nil}^*(\mathbb{Z}_{p^2}))) \cong S_{p-1}$. This confirms that the decomposition theorem of the preceding section applies nontrivially only for $k \geq 3$, while $k = 2$ represents the unique complete-graph exception.

7 Conclusion

This paper provides a complete and explicit determination of the automorphism groups of zero-divisor graphs arising from the nonzero nilradical of finite local rings of the form \mathbb{Z}_{p^k} . The central mechanism underlying these results is the stratification of nilpotent elements by p -adic valuation, which induces a rigid layered structure on the vertex set of the graph. These valuation layers serve as invariant combinatorial units: vertices within a single layer are indistinguishable, while vertices across layers are separated by degree and adjacency constraints. As a consequence, the automorphism group decomposes naturally as a direct product of symmetric groups acting independently on each valuation level. This valuation-driven perspective not only clarifies the symmetry structure of $\Gamma(\text{Nil}^*(\mathbb{Z}_{p^k}))$ but also establishes a foundational case for broader investigations. In subsequent work, similar techniques can be adapted to semilocal rings and mixed-exponent settings, where multiple valuation systems interact to shape graph automorphisms in more intricate ways.

Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Competing Interests

Authors have declared that no competing interests exist.

References

- Akbari, S., Mohamadian, H., and Yassemi, S. (2003). When a zero-divisor graph is planar or a complete r -partite graph. *J. Algebra*, 270:169–180.
- Ali, N., Siddiqui, H. M. A., and Qureshi, M. I. (2025). Characterizing rings based on resolvability in associated compressed zero divisor graphs. *J. Algebra Appl.*, page 2541004.
- Ali, N., Siddiqui, H. M. A., Riaz, M. B., Qureshi, M. I., and Akgül, A. (2024). A graph-theoretic approach to ring analysis: dominant metric dimensions in zero-divisor graphs. *Heliyon*, 10(10).
- Anderson, D. F. and Badawi, A. (2008). On the zero-divisor graph of a ring. *Commun. Algebra*, 36:3073–3092.
- Anderson, D. F. and Livingston, P. S. (1999). The zero divisor graph of a commutative ring. *J. Algebra*, 217:437–447.
- Axtell, M. and Stickles, J. (2009). Zero divisor ideals and realizable zero divisor graphs. *J. Math.*, 2.
- Beck, I. (1988). Coloring of commutative rings. *J. Algebra*, 116:208–226.
- DeMeyer, F. and Schneider, K. (2003). Automorphisms and zero divisor graphs of commutative rings. In Badawi, A., editor, *Trends in commutative rings research*, pages 93–105. Nova Science Publishers.

- Lucas, T. G. (2006). The diameter of a zero divisor graph. *J. Algebra*, 301:174–193.
- Madhumitha, S. and Naduvath, S. (2023). Graphs defined on rings: a review. *Mathematics*, 11(17):3643.
- Mude, L. H., Oduor, O. M., and Onyango, O. M. (2023). Automorphisms of zero divisor graphs of power four radical zero completely primary finite rings. *Asian Res. J. Math.*, 19(8):108–113.
- Onyango, O. M., Oduor, O. M., and Odhiambo, P. O. (2016). Automorphisms of the unit groups of square radical zero finite commutative completely primary rings. *Int. J. Pure Appl. Math.*, 108(1):39–48.
- Patil, A., Khairnar, A., and Momale, P. S. (2022). Zero-divisor graph of a ring with respect to an automorphism. *Soft Comput.*, 26(5):2107–2119.
- Sbarra, E. and Zanardo, M. (2024). Zero-divisor graphs and zero-divisor functors. *J. Algebra Appl.*, 23(12):2450199.
- Spiroff, S. and Wickham, C. (2011). A zero divisor graph determined by equivalence classes of zero divisors. *Commun. Algebra*, 39:2338–2348.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2026): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)
<https://pr.sdiarticle5.com/review-history/152768>